

## Getting the most from Grazon Pro

- Read the label before use
- Only apply this product if you have PA1 and PA6 certificates or similar qualifications (see Key Points)
- Ensure that your Knapsack Sprayer is serviceable and calibrated prior to use
- Wear suitable PPE at all times during the preparation, spraying and cleaning process
- For optimum results spray weeds whilst actively growing and before they flower, or top them first and spray regrowth after 2-3 weeks
- If weeds are too tall or too dense, 'shadowing' can occur so, again, top them first and spray regrowth after 2-3 weeks
- Not all weeds will be at the best growth stage for spraying at the time of application, so a follow-up treatment may be necessary
- Spray weed leaves until thoroughly wet but avoid run-off
- Always direct spray away from water
- Other people and pets should keep out of treated areas until the spray has dried on treated foliage
- Re-seed bare patches after weeds have died to avoid encroachment by new weeds.

### Treat target weeds at optimal size.

If they are too big then top and spray regrowth. This photo shows the target weed at optimal size.



**Grazon® Pro** – is ideal for spot treatment of a wide range of difficult to control weeds. Use on small patches, isolated weeds, on steep ground or under fence lines. It controls docks, thistles, nettles, brambles, gorse and broom.

**See product label for full details.**

For use **ONLY** with a knapsack or hand-held lance



# Grazon® PRO

## HERBICIDE

# Spot treatment for docks, nettles and thistles.



## Forage knowledge on the go.



Free and easy to use, the Corteva Forage App is packed with information and tools to help you maximise your grass and maize crops. Simply scan the QR code.



For grassland advice call Whelehan Crop Protection on: **01 574 2410** or visit: [www.corteva.ie/forage](http://www.corteva.ie/forage) or email: [CropProtection@tpwhelehan.ie](mailto:CropProtection@tpwhelehan.ie)  
Follow us: [X@CortevaForage](https://twitter.com/CortevaForage) or [@CortevaIE](https://www.instagram.com/CortevaIE)

**Whelehan**  
CROP PROTECTION

**CORTEVA**  
agriscience

Use plant protection products safely. Always read the label and product information before use. For further information including warning phrases and symbols refer to label. Triple rinse containers, puncture and invert to dry at time of use.

Authorisation Holder: Corteva Agriscience UK Limited, CPC2 Capital Park, Fulbourn, Cambridge CB21 5XE. Tel: +44 1462 457272. Marketing Company: Whelehan Crop Protection, Suite 11/12 Bunkilla Plaza, Bracetown Business Park, Clonee, Dublin 15, Co Dublin Ireland. Tel: 01 574 2410 Email: [cropprotection@tpwhelehan.ie](mailto:cropprotection@tpwhelehan.ie) \* Trademarks of Corteva Agriscience and its affiliated companies. Grazon® Pro contains clopyralid and triclopyr.  
©2024 Corteva. Supercedes all previous editions of this leaflet.

**CORTEVA**  
agriscience

## HERBICIDE

The ideal spot treatment solution for a wide range of difficult to control weeds where safety to grass is important.

### Use GrazonPro because it is:

- Approved for Knapsack Sprayer use in established grassland
- Fast-acting and controls a wide range of broad-leaved and 'woody' weeds – see 'Weeds controlled' table
- Offers excellent grass safety
- Rainfast in 2 hours
- Needs only 7-day post-treatment stock exclusion interval\*
- Economic to use;
  - low dose rate of just 60ml per 10L water
  - 10 fills of a 16lt Knapsack Sprayer from a single 1lt pack
  - it comes with a FREE measuring beaker.

\* In the absence of Ragwort



### Key points:

<b>Active ingredients</b>	240g/l triclopyr + 60g/l clopyralid
<b>'For Professional Use'</b>	Grazon Pro is For Professional Use.
<b>Weeds controlled</b>	   <p>Docks      Nettles      Thistles</p>
<b>Application rate</b>	60ml Grazon Pro per 10 litres water
<b>Application method</b>	Knapsack Sprayer or other hand-held lance
<b>Treatment timing</b>	Between 1st March and 31st October
<b>Weed size</b>	
<b>Broadleaved dock and Curled dock</b>	Rosette stage, 150mm to 250mm across or high
<b>Common nettle</b>	Actively growing before flowering
<b>Creeping thistle and Spear thistle</b>	Rosette stage, flowering spike up to 150mm high
<b>Bramble, Broom and Gorse</b>	Between June and August when actively growing, before the onset of senescence. Foliage must be thoroughly wetted.
<b>Weed health</b>	Weeds must be actively growing; free from disease and insect damage; not suffering from drought, waterlogging or nutrient deficiency
<b>Stock exclusion</b>	Keep livestock out of treated areas for at least 7 days*
<b>Cutting interval (pre-treatment)</b>	Leave 14 - 21 days to allow sufficient regrowth of both grass and weeds
<b>Cutting interval (post-treatment)</b>	To allow maximum translocation of GrazonPro to the weed roots, do not cut grass for 28 days
<b>Rainfastness</b>	2 hours when applied in dry conditions to a dry leaf
<b>Clover</b>	GrazonPro will damage clover, but a well-aimed spray onto the target weeds will enable high levels of weed control to be achieved whilst minimizing the effect on the overall clover population

\* In the absence of Ragwort

### \*Ragwort label guidance

Ragwort plants sprayed with this herbicide are more palatable and contain higher levels of toxins. Animals should be excluded from treated areas until any ragwort has completely recovered or died and there is no visible sign of the dead weed. Do not include treated ragwort in hay or silage crops.



## Weeds controlled

Where we have knowledge of how our products might affect other grassland weeds we have detailed it in the following tables. These are not recommendations; just an indication of what control might be achieved.   
 ▽ indicates information based on anecdotal or limited data, and as such the user bears the risk in respect of failures concerning efficacy and phytotoxicity.

### Annual weeds

Bindweed (black)	Good control
Bristly ox-tongue	Moderate control
Charlock	Good control
Chickweed	Good control
Cleavers	Good control
Corn chamomile	Good control
Corn marigold	Good control
Cranesbill	Good control
Dead-nettles	Good control
Fat-hen	Good control
Fool's parsley	Good control
Forget-me-not	Good control
Fumitory	Good control
Groundsel	Good control
Hemp-nettle	Good control
Himalayan balsam	Good control
Knotgrass	Good control
Mayweeds	Good control
Nettle (small)	Good control
Nightshade (black)	Good control
Orache	Good control
Pale persicaria	Good control
Poppy	Good control
Redshank	Good control
Scarlet pimpernel	Moderate control
Shepherd's-purse	Good control
Speedwells	Moderate control
Spurrey	Good control
Wild radish	Good control
Yellow rattle	Moderate control

### Perennial weeds

Bindweed (field)	Good control
Bracken	Good control
Bramble	Good control
Broom	Good control
Burdock	Good control
Buttercups	Some control
Cinquefoil (creeping)	Good control
Clover, trefoil	Good control
Coltsfoot	Good control
Cow parsley	Moderate control
Daisy (common)	Moderate control
Daisy (ox-eye)	Good control
Dandelion	Moderate control
Docks	Good control
Gorse	Good control
Ground elder	Good control
Ground ivy	Good control
Hawthorn	Good control
Hemlock	Good control
Hogweed	Good control
Horsetail ( <i>Equisetum</i> )	Some control
Japanese knotweed	Good control
Knapweed (common)	Moderate control
Lesser celandine	Good control
Mallow (common)	Good control
Medick (black)	Good control
Mugwort (common)	Good control
Nettle (common)	Good control
Old man's beard	Good control
Plantain (greater)	Moderate control
Plantain (ribwort)	Moderate control
Ragwort	Some control
Rosebay willowherb	Good control
Rushes	Moderate control
Self-heal	Good control
Silverweed	Good control
Sorrel (common)	Good control
Thistles	Good control
Vetch, tare	Good control
Yarrow	Moderate control
Yellow/Flag Iris	Good control

### Weed control key

Good control
Moderate control
Some control
No control
No information
Anecdotal or limited information